

Irreducible components of the equivariant punctual Hilbert schemes

Laurent Evain (evain@tonton.univ-angers.fr)

June 10, 2001

Abstract: Let $\mathbb{H}_{ab}$ be the equivariant Hilbert scheme parametrizing the 0-dimensional subschemes of the affine plane invariant under the natural action of the one-dimensional torus $T_{ab} := \{(t^{-b}, t^a) \mid t \in k^*\}$. We compute the irreducible components of $\mathbb{H}_{ab}$: they are in one-one correspondence with a set of Hilbert functions.

As a by-product of the proof, we give new proofs of results by Ellingsrud and Strømme, namely the main lemma of the computation of the Betti numbers of the Hilbert scheme $\mathbb{H}^l$ parametrizing the 0-dimensional subschemes of the affine plane of length l [4], and a description of Bialynicki-Birula cells on $\mathbb{H}^l$ by means of explicit flat families [5]. In particular, we precise conditions of applications of this last description.

Introduction

The Hilbert scheme $\mathbb{H}$ parametrizing subschemes of the plane $\text{Spec } k[x, y]$ is a disjoint union of its components $\mathbb{H}^l$ parametrizing the subschemes of length l . Two auxiliary subschemes are useful to study $\mathbb{H}$. The first one is $\mathbb{H}_*$ parametrizing the subschemes supported by a point $*$. For instance, the papers by Lehn and Nakajima ([12] and [13]) are explicit situations where the geometry of $\mathbb{H}_*$ has proven to be useful. The second auxiliary scheme $\mathbb{H}_{ab}$ encode in some sense the action of the torus on $\mathbb{H}$. The action we are talking about is the action of the two dimensional torus $k^* \times k^*$ induced by the linear action $(t_1, t_2).x^\alpha y^\beta = (t_1.x)^\alpha (t_2.y)^\beta$ of $k^* \times k^*$ on $k[x, y]$. If $T_{ab} = \{(t^{-b}, t^a) \mid t \in k^*\}$ is a one dimensional sub-torus, the closed subscheme $\mathbb{H}_{ab} \subset \mathbb{H}$ parametrizes by definition the zero dimensional subschemes invariant under the action of T_{ab} . Its usefulness has been illustrated by Ellingsrud and Strømme's work [4] where they compute the Chow group of $\mathbb{H}^l$ by studying the equivariant embedding $\mathbb{H}_{ab} \hookrightarrow \mathbb{H}$ for (a, b) general enough. See Brion [3] for other links between $\mathbb{H}$ and $\mathbb{H}_{ab}$ relative to equivariant cohomology. So, even if one is primarily interested in studying $\mathbb{H}$, it is natural to study the three Hilbert schemes $\mathbb{H}$, $\mathbb{H}_{ab}$ and $\mathbb{H}_*$ as a whole for their respective geometries are linked.

One of the basic questions about these schemes is to determine their irreducible components. The answers are known for $\mathbb{H}$ (the irreducible components are the smooth subschemes $\mathbb{H}^l$ by Hartshorne [9] and Fogarty [7]) and for $\mathbb{H}_*$ (the irreducible components are $\mathbb{H}_*^l := \mathbb{H}_* \cap \mathbb{H}^l$ by Briançon [2]), but unknown for $\mathbb{H}_{ab}$.

As to $\mathbb{H}_{ab}$, it cannot be irreducible for there are obviously disconnected subschemes $\mathbb{H}_{ab}^l := \mathbb{H}_{ab} \cap \mathbb{H}^l$. These can still be divided into smaller disconnected pieces as follows. The ideal of a subscheme corresponding to a point of $\mathbb{H}_{ab}^l$ is quasi-homogeneous (homogeneous if $a = 1, b = -1$) and is a direct sum $I = \oplus_{l \in \mathbb{Z}} I_l$ of vector spaces. The codimensions d_i of the I_i are fixed on a connected component and verify $\sum d_i = l$. Thus $\mathbb{H}_{ab}^l = \coprod H_{ab}(H)$, where $H : \mathbb{Z} \rightarrow \mathbb{N}$ runs through the functions verifying $\sum_{i \in \mathbb{Z}} H(i) = l$ and $H_{ab}(H)$ parametrizes the zero-dimensional subschemes satisfying $(\dots, d_{-1}, d_0, d_1, d_2, \dots) = H$. So the natural question concerning the irreducibility of the equivariant Hilbert schemes is: are the natural candidates $\mathbb{H}_{ab}(H)$ connected and irreducible?

Theorem 1. *For any H , if $\mathbb{H}_{ab}(H) \neq \emptyset$, then $\mathbb{H}_{ab}(H)$ is smooth and connected (hence irreducible).*

This result was already known for (a, b) general enough (since then $\mathbb{H}_{ab}^l$ is a union of points) and for the special value $(a, b) = (1, -1)$ thanks to Iarrobino [10].

On the proof.

The first point is that $\mathbb{H}_{ab}$ is smooth as it is the fixed locus of the smooth scheme $\mathbb{H}$ under the action of the one-dimensional torus T_{ab} [11], so we just have to prove the connectedness of $\mathbb{H}_{ab}(H)$.

The general plan is as follows. We explain that $\mathbb{H}_{ab}(H)$ is naturally stratified by locally closed subschemes $C(E)$. The strata are parametrized by staircases (or Young diagrams) E and in each stratum $C(E)$ there is a particular subscheme $Z(E)$. Each stratum is connected as an affine space, so the problem is to link the strata together. This is done in two steps. Firstly, we partially order the strata and we show that each stratum is connected to a smaller stratum except for some special strata characterized by their staircase E . We then show that there is in fact a unique special stratum. It follows that all strata can be connected to this special stratum and the theorem follows. The main point is to find a workable condition to connect a stratum $C(E)$ to a smaller stratum. To this purpose, we describe the tangent space $T_{Z(E)}^{ab}$ to $\mathbb{H}_{ab}(H)$ at the special point $Z(E)$ of the stratum. It is a vector space whose a base is a combinatorial datum associated with E (theorem 3). This vector space splits into a direct sum of a positive tangent space $T_{Z(E)}^{ab+}$ and a negative tangent space $T_{Z(E)}^{ab-}$. We describe a sort of “exponential” map $T_{Z(E)}^{ab+} \rightarrow \mathbb{H}_{ab}$, which turns out to be an isomorphism on $C(E)$. Using this map, a point p of $T_{Z(E)}^{ab+}$ corresponds to a subscheme $X(p)$ of $\mathbb{A}^2$ whose equations are completely described. We check that for a suitable action of k^* , if $p \neq 0$, then $\lim_{t \rightarrow 0} t.X(p)$ is a subscheme in a smaller stratum (proposition 18). This allows us to connect a stratum $C(E)$ to a smaller stratum provided the positive tangent space in $Z(E)$ is non trivial.

To conclude, we exhibit an explicit staircase E_m characterized combinatorially by a property of minimality (theorem 19). We show that going down from a stratum to a smaller stratum as explained above, the process always ends in $C(E_m)$. The existence of a minimal E_m is a purely combinatorial statement. Its proof, however, requires the characterization of minimality using the positive tangent space. The method here is

very much in the spirit of toric varieties [8] where combinatorial statements are proved via algebro-geometric arguments.

By-products of the proof. The methods involved in the proof to study $\mathbb{H}_{ab}$ remain useful with few changes to study $\mathbb{H}$. This allows us to recover results by Ellingsrud-Strømme ([4] and [5]).

More precisely, the tangent space $T_{Z(E)}$ to $\mathbb{H}$ at $Z(E)$ is a direct sum of the various invariant tangent spaces $T_{Z(E)}^{ab}$. Following the lines of [4], the control of $T_{Z(E)}$ is the key to compute the Betti numbers of $\mathbb{H}^l$ and we get by our description of $T_{Z(E)}$ a new proof of their main technical lemma.

This tangent space $T_{Z(E)}$ splits into a direct sum $T_{Z(E)}^+ \oplus T_{Z(E)}^-$, just like the invariant tangent space $T_{Z(E)}^{ab}$. And we can still define an embedding $e : T_{Z(E)}^+ \rightarrow \mathbb{H}$ using the same methods as for the invariant case, that is by exhibiting an explicit flat family on $T_{Z(E)}^+$. The image of e is a Bialynicki-Birula cell $C(E)$ with respect to a suitable action of the torus k^* on $\mathbb{H}$ (theorem 13). In other words, we have explicit charts for cells $C(E)$. Though it has been obtained by quite different methods, the flat family which defines e is up to an identification the same family as in [5]. We note that the families involved in [5] are more general than ours since they give explicit charts for cells $C(E)$ defined by a more general action of the torus k^* . However, some of the extra cases may have gaps. Consequently, theorem 13 can be seen as a prolongation of the analysis of Ellingsrud and Strømme: it provides conditions on the action of k^* under which the description of the cells given in [5] are valid (section 3.2).

1 The stratified subscheme $\mathbb{H}_{ab}(H)$.

1.1 Disconnected subschemes of $\mathbb{H}_{ab}$

In this section, we introduce the closed subschemes $\mathbb{H}_{ab}(H) \subset \mathbb{H}_{ab}$. As claimed in the introduction, for $H_1 \neq H_2$, $\mathbb{H}_{ab}(H_1) \cap \mathbb{H}_{ab}(H_2) = \emptyset$.

Recall that k is an algebraically closed field, that a and b are two relatively prime integers, and that $T_{ab} = \{(t^{-b}, t^a), t \in k^*\}$ is a one dimensional subtorus of $k^* \times k^* =: T$. Since $(a, b) \neq (0, 0)$ we can suppose by symmetry $b \neq 0$, and since (a, b) is defined up to sign, we suppose from now on $b < 0$. The Hilbert scheme $\mathbb{H}^l$ parametrizing the zero-dimensional subschemes of $\text{Spec } k[x, y]$ of length l is connected and by definition $\mathbb{H} = \amalg \mathbb{H}^l$ where $\amalg$ stands for a disjoint union. There are actions $T \times \mathbb{H} \rightarrow \mathbb{H}$ and $T_{ab} \times \mathbb{H} \rightarrow \mathbb{H}$ of the tori on $\mathbb{H}$, which in terms of coordinates are given by $(t_1, t_2) \cdot (x^\alpha y^\beta) = (t_1 x)^\alpha (t_2 y)^\beta$. We denote by $\mathbb{H}_{ab} \subset \mathbb{H}$ the subscheme parametrizing the subschemes of the plane invariant under the action of T_{ab} . Then $\mathbb{H}_{ab} = \amalg \mathbb{H}_{ab}^l$, where $\mathbb{H}_{ab}^l := \mathbb{H}_{ab} \cap \mathbb{H}^l$. One can still separate $\mathbb{H}_{ab}^l$ into disjoint subschemes by fixing a Hilbert function.

To do this, let us characterize the subschemes of $\text{Spec } k[x, y]$ which lie in $\mathbb{H}_{ab}$ by their ideals. Define the degree d of a monomial by the formula $d(x^\alpha y^\beta) = -b\alpha + a\beta$. If I

is an ideal of $k[x, y]$, we let $I_n := I \cap k[x, y]_n$, where $k[x, y]_n$ denotes the vector space generated by the monomials m of degree $d(m) = n$. A subscheme Z is in $\mathbb{H}_{ab}$, if and only if its ideal is quasi-homogeneous with respect to d , ie. $I(Z) = \bigoplus_{n \in \mathbb{Z}} I(Z)_n$.

By semi-continuity, if a subscheme $Z' \in \mathbb{H}_{ab}$ is a specialization of $Z \in \mathbb{H}_{ab}$, then the codimensions of $I_n(Z)$ and $I_n(Z')$ in $k[x, y]_n$ verify $\text{codim} I_n(Z) \geq \text{codim} I_n(Z')$. But Z and Z' , being in the same connected component of $\mathbb{H}$, have the same length $l = \sum_n \text{codim} I_n(Z) = \sum_n \text{codim} I_n(Z')$. It follows that the sequence $H(Z) = (\dots, h_{-1}, h_0, h_1, \dots)$, where $h_i = \text{codim} I_n(Z)$, is constant on the connected components of $\mathbb{H}_{ab}$ and that $h_n = 0$ for $n \gg 0$ and $n \ll 0$. If $H = (\dots, 0, 0, h_r, \dots, h_s, 0, 0, \dots)$ is any sequence, we denote by $\mathbb{H}_{ab}(H)$ the closed subscheme of $\mathbb{H}_{ab}$ (possibly empty) parametrizing the subschemes Z verifying $H(Z) = H$. By the above, we have $\mathbb{H}_{ab} = \coprod \mathbb{H}_{ab}(H)$.

1.2 The case $ab \geq 0$ via a theorem of Bialynicki-Birula

In this section, we introduce the stratification on $\mathbb{H}_{ab}(H)$, the special points $Z(E)$ of the cells $C(E)$ and we prove the theorem under the condition $ab \geq 0$. The strata can be defined in terms of Gröbner bases, using Grassmannians or using a Bialynicki-Birula theorem, as explained in [6]. We recall here the first and the last approach. The theorem follows easily if $ab \geq 0$: the stratification on $\mathbb{H}_{ab}(H)$ is reduced to one stratum, which is an affine space by Bialynicki-Birula.

Each cell is associated with a staircase E : to a staircase E corresponds a subscheme $Z(E)$ and the cell $C(E)$ is the unique cell containing the subscheme $Z(E)$. To be more precise, we recall that a staircase is a subset of $\mathbb{N}^2$ whose complementary is stable by addition of $\mathbb{N}^2$. In this paper, we will identify freely the monomial $x^p y^q$ with the couple (p, q) and therefore the expression “staircase of monomials” will make sense. More generally, we will transpose unscrupulously the definitions between couples of integers and monomials. If E is a staircase, then the vector space I^E generated by the monomials which are not in E is an ideal and conversely, every monomial ideal is an ideal I^E for a unique staircase E . The subscheme $Z(E)$ whose ideal is I^E is in $\mathbb{H}_{ab}(H)$ if and only if E has $H(i)$ elements in degree i .

The Gröbner bases point of view. We choose to order the monomials of $k[x, y]$ by the rule: $m_1 < m_2$ if $(d(m_1), d_y(m_1)) < (d(m_2), d_y(m_2))$ for the lexicographic order, where d_y is defined by $d_y(x^\alpha y^\beta) = \beta$. Let $m_1, m_2, \dots$ be the monomials which don't belong to E . Fixing once and for all a, b and $H = (\dots, 0, 0, h_r, \dots, h_s, 0, \dots)$, an ideal of $\mathbb{H}_{ab}(H)$ is in $C(E)$ if, regarding it as a k -vector space, it admits a basis $f_1, f_2, \dots$ where $f_i = m_i + R_i$, R_i being a linear combination of monomials strictly smaller than m_i . The locus $C(E)$ in $\mathbb{H}_{ab}$ is non empty exactly when E has h_i elements in degree i . If $ab < 0$, the above order is a monomial order in the sense of Gröbner bases. In this case, the theory of Gröbner bases associates with every ideal in $k[x, y]$ a monomial ideal called initial ideal and $C(E)$ is the locus in $\mathbb{H}_{ab}(H)$ parametrizing the ideals whose initial ideal is I^E .

The Bialynicki-Birula point of view. Let X be a smooth projective variety over k admitting an action of the torus k^* . Suppose that the action has a finite number of fixed points $x_1, \dots, x_n$. Let T_{X, x_i}^+ be the part of the tangent space to x_i in X where the weights of the k^* -action are positive, and let $X_i := \{x \in X, \lim_{t \rightarrow 0}(t.x) = x_i\}$. Then a theorem of Bialynicki-Birula asserts that the X_i are a cellular decomposition of X in affine spaces and satisfy $T_{X_i, x_i} = T_{X, x_i}^+$. If $U \subset X$ is a stable open subset of X , it is still possible to define cells which are affine spaces associated with the fixed points which lie in U , but these cells do not always cover U . In our case, fixing two integers p and q with $ap + bq > 0$, the torus k^* acts on $k[x, y]$ by $t.x = t^p x$ and $t.y = t^q y$. This action induces an action of k^* on $\mathbb{H}_{ab}(H)$. The fixed points of $\mathbb{H}_{ab}(H)$ under k^* are the monomial subschemes $Z(E)$. Applying the Bialynicki-Birula theorem to the action of k^* on $\mathbb{H}_{ab}(H)$, we get a set of cells. We denote by $C(E)$ the cell associated with the fixed point $Z(E)$. The previous description of these cells insure that they cover $\mathbb{H}_{ab}(H)$.

Proof of the theorem in the case $ab \geq 0$. When the product ab is non negative, there is at most one staircase E compatible with the Hilbert function H (i.e. such that E has h_i elements in degree i). It follows that $\mathbb{H}_{ab}(H)$ is empty or an affine space $C(E)$. The theorem is then obvious.

2 Description of the tangent space

Let E be a staircase, $T_{Z(E)}$ be the tangent space to $\mathbb{H}$ at the point $Z(E)$, $T_{Z(E)}^{ab}$ be the tangent space to $\mathbb{H}_{ab}$ at the point $Z(E)$. In this section, we give a description of $T_{Z(E)}^{ab}$ and $T_{Z(E)}$.

We need some combinatorial vocabulary that we introduce now.

Clefts and cleft couples. A cleft for E is a couple (u, v) such that $x^u y^v$ is a monomial in I^E minimal for the divisibility relation among the monomials of I^E . A positive (resp. negative) half-direction is a couple of relatively prime integers (f, g) with $f > 0$ (resp. $f < 0$) or $f = 0$ and $g < 0$ (resp. $g > 0$). A couple of points (M, N) in $\mathbb{N}^2$ has half-direction (f, g) if the vector $\vec{MN}$ is a positive multiple of (f, g) . We have a notion of direction by identifying two opposites half directions. A cleft couple (relatively to E) with half-direction h is a couple of elements (c, m) in $\mathbb{N}^2$ s.t. c is a cleft, $m \in E$, and (c, m) has half-direction h .

The orders $>_+$ and $>_-$ on cleft couples. We put orders on the set of clefts and on the set of cleft couples. The identification between $x^p y^q$ and (p, q) gives a lexicographic order $>_+$ on the monomials of $k[x, y]$. The identification between $x^p y^q$ and (q, p) gives a lexicographic order $>_-$. Any order on the monomials induce an order on the clefts (by restriction), on the couples of monomials (lexicographically) and on the cleft couples (the restriction of the latter). We still note the induced orders $>_+$ and $>_-$.

Some particular sets of cleft couples and associated vector spaces. The combinatorial object used to describe $T_{Z(E)}$ is the significant cleft couple:

Definition 2. Let (c, m) be a cleft couple with positive (resp. negative) half-direction, c' be the cleft successor of c for $>_+$ (resp. for $>_-$) and s be the smaller common multiple of c and c' . Then (c, m) is said to be a significant couple if $m \frac{s}{c} \notin E$.

For a cleft couple, we'll be interested to know whether it is significant and what is its direction. Formally, we introduce the sets $C, \overline{C}, C_{ab}, \overline{C}_{ab}$ containing respectively

- the significant cleft couples (with respect to E)
- the cleft couples
- the significant cleft couples with direction (a, b) .
- the cleft couples with direction (a, b)

We have the obvious inclusions:

$$\begin{array}{ccc} C & \rightarrow & \overline{C} \\ \uparrow & & \uparrow \\ C_{ab} & \rightarrow & \overline{C}_{ab} \end{array}$$

For each of the above sets, we can form the vector space on this set, i.e. the vector space whose elements are the formal linear combinations of elements of this set. We denote by $R, \overline{R}, R_{ab}, \overline{R}_{ab}$ the vector spaces corresponding to $C, \overline{C}, C_{ab}, \overline{C}_{ab}$. To the inclusion of sets corresponds the inclusion of vector spaces

$$\begin{array}{ccc} R & \rightarrow & \overline{R} \\ \uparrow & & \uparrow \\ R_{ab} & \rightarrow & \overline{R}_{ab} \end{array}$$

We can now formulate the main result of the section, whose proof ends with corollary 6.

Theorem 3. *There is a natural isomorphism $T_{Z(E)}^{ab} \simeq R_{ab}$.*

We start by constructing an injective morphism $\varphi : T_{Z(E)}^{ab} \rightarrow \overline{R}$. Next, we will identify the image $\varphi(T_{Z(E)}^{ab})$ with R_{ab} .

We have the classical description of tangent spaces on Hilbert schemes $T_{Z(E)} \simeq \text{Hom}_{k[x,y]}(I^E, k[x,y]/I^E)$. Let's recall in our situation how to produce the infinitesimal deformation of $k[x,y]/I^E$ starting from an element $f \in \text{Hom}_{k[x,y]}(I^E, k[x,y]/I^E)$. Let V_E be the vector space generated by the monomials which are in E , $\pi : V_E \rightarrow k[x,y]/I^E$ the isomorphism induced by the restriction of the projection $k[x,y] \rightarrow k[x,y]/I^E$, and $p : k[x,y]/I^E \rightarrow k[x,y]$ the inverse of π . Let $\overline{f} = p \circ f : I^E \rightarrow k[x,y]$. The set of elements $J = \{m + \epsilon \overline{f}(m), m \in I^E\}$ is an ideal of $k[x,y][\epsilon]/(\epsilon^2)$. The quotient $k[x,y][\epsilon]/J$ is the flat infinitesimal deformation of $k[x,y]/I^E$ corresponding to f . For every cleft couple (c, m) , define $\lambda_{c,m}$ by the formula $f(c) = \sum \lambda_{c,m} m$.

Proposition 4. *The linear map*

$$\begin{array}{ccc} \varphi : & T_{Z(E)} & \rightarrow \overline{R} \\ f \text{ s.t. } \overline{f}(c) = \sum \lambda_{c,m} m & \mapsto & \sum \lambda_{c,m}(c, m) \end{array}$$

is injective and its restriction to $T_{Z(E)}^{ab}$ factorizes:

$$\begin{array}{ccc} T_{Z(E)}^{ab} & \rightarrow & \overline{R} \\ & \searrow & \uparrow \\ & & \overline{R}_{ab} \end{array}$$

Proof: the ideal I^E being generated by the clefts, the morphism $f \in \text{Hom}_{k[x,y]}(I^E, k[x,y]/I^E)$ is characterized by the images of the clefts. But these images are themselves characterized by $\varphi(f)$ so φ is injective.

By Bialynicki-Birula [1], $T_{Z(E)}^{ab} \subset T_{Z(E)}$ is the subspace containing the vectors fixed under the action of T_{ab} . If f is in $T_{Z(E)}^{ab} \subset T_{Z(E)}$, one then sees from the description of f as an infinitesimal deformation of $k[x,y]/I$ that $\lambda_{c,m}$ is different from zero only if (c, m) has direction (a, b) . It follows that $\varphi(T_{Z(E)}^{ab}) \subset \overline{R}_{ab}$. ■

The next step is to show the equality $\varphi(T_{Z(E)}^{ab}) = R_{ab} \subset \overline{R}_{ab}$. This is done in two steps. Firstly, we construct a graph G associated to E , a vector space R_G from G and we show the equality $\varphi(T_{Z(E)}^{ab}) = R_G$. We then conclude with the isomorphism $R_G \simeq R_{ab}$.

We construct a graph G from the staircase E . The set of points is the set C_{ab} of cleft couples with direction (a, b) . Significant couples are not the end of any arrow. Non significant cleft couples are the end of exactly one arrow. Let (c_1, m_1) be non significant with positive half direction, and c_2 be the successor cleft of c_1 for $>_+$. If $m_2 := m_1 \cdot \frac{c_2}{c_1} \in k[x, y]$, draw an arrow from (c_2, m_2) to (c_1, m_1) , otherwise draw an arrow from (c_1, m_1) to itself. Replacing $>_+$ by $>_-$, you get arrows ending at cleft couples with negative half-direction.

If G is a graph with set of points $\mathcal{P} = \{p_1, \dots, p_n\}$ and arrows $f_1, \dots, f_p$, denote $o(f_i)$ the origin of f_i and $e(f_i)$ the end of f_i . Let $R_{\mathcal{P}}$ be the vector space whose base is the set of elements of $\mathcal{P}$ and R_G be the sub-vector space of $R_{\mathcal{P}}$ whose elements are the linear combinations $\sum \lambda_{p_i} p_i$ verifying $\lambda_{p_i} = \lambda_{p_j}$ if there is an arrow from p_i to p_j and $\lambda_{p_i} = 0$ if there is an arrow from p_i to itself.

Proposition 5. *If $a.b \leq 0$, then $\varphi(T_{Z(E)}^{ab}) = R(G)$. If $a.b > 0$, then $\varphi(T_{Z(E)}^{ab})$ is a subspace of $R(G)$.*

Proof: we have seen that an element f in $T_{Z(E)} = \text{Hom}_{k[x,y]}(I^E, k[x,y]/I^E)$ is characterized by the images $g_i = f(c_i)$ of the clefts c_i . Reciprocally, if we prescribe an image g_i for each cleft c_i , a compatibility relation insures the existence of f in $T_{Z(E)}$ sending c_i to g_i : if $c_i <_+ c_j$ are two clefts such that c_j is the successor of c_i , and if s is the smaller common multiple (s.c.m.) of c_i and c_j , then their images g_i, g_j have to verify

$$g_i \cdot \frac{s}{c_i} = g_j \cdot \frac{s}{c_j}$$

The element g_i is a linear combination $\sum \lambda_{i,k} \overline{m_k}$ and we identify it with the element $\sum \lambda_{i,k}(c_i, m_k)$ using φ . We do similarly for g_j . The above relation between g_i and g_j translates into relations between the coefficients $\lambda_{i,k}$ and $\lambda_{j,l}$. One sees that these relations are the relations given by the arrows of the graph in the case $ab \leq 0$ so we are done. In the case $ab > 0$, there are at least these relations and possibly some others. ■

Corollary 6. *If $a.b \leq 0$, then $T_{Z(E)}^{ab} \simeq R_{ab}$. If $a.b > 0$, then $T_{Z(E)}^{ab} = R_{ab} = 0$.*

Proof: by construction, each point in the graph G is the end of at most one arrow and the origin of at most one arrow. It follows that the connected components of the graph G are chains of consecutive points $p_1, p_2, \dots, p_n$ where two points p_i and p_{i+1} are connected by one arrow from p_i to p_{i+1} . Moreover the p_i are distinct if $n \geq 2$. The vector space $R(G)$ is then obviously isomorphic to $R(G')$, where G' is the graph with no arrow obtained from G by keeping the points that are not the end of any arrow. The set of points of G' is just the set C_{ab} of significant cleft couples with direction (a, b) so $R(G) = R(G') = R_{ab}$ and the first claim of the corollary is a consequence of the previous proposition. In the case $a.b > 0$, G' is empty because there are no significant cleft couples. So $R(G) = 0$ and the second claim follows again from the last proposition. ■

This corollary obviously concludes the proof of theorem 3. It also clarifies the structure of $\mathbb{H}_{ab}$ in some cases. We have already explained that $\mathbb{H}_{ab}$ is a disjoint union of affine spaces if $ab \geq 0$. These affine spaces are particularly simple if $ab > 0$:

Corollary 7. *If $ab > 0$, then $\mathbb{H}_{ab}(H)$ is empty or reduced to a point.*

Proof: there is at most one staircase E compatible with H . If E does not exist, then $\mathbb{H}_{ab}(H) = \emptyset$. Otherwise, $\mathbb{H}_{ab}(H) = C(E)$. The cell $C(E)$ is an affine space by Bialynicki-Birula and its dimension is zero by the previous corollary (6) ■

Following the lines of the proof of theorem 3 and simply forgetting the arguments concerning the action of the torus T_{ab} , we have a description of the tangent space $T_{Z(E)}$ instead of a description of the invariant tangent space $T_{Z(E)}^{ab}$.

Theorem 8. *The tangent space $T_{Z(E)}$ to $\mathbb{H}$ at $Z(E)$ is isomorphic to the vector space R whose base is the set of significant cleft couples of E .*

Remark 9. *This description of the tangent space implies the main lemma (3.2) of [4]*

2.1 Definition of the positive and negative tangent spaces

Let C_{ab+} (resp. C_{ab-}) be the set of significant cleft couples with direction (a, b) and positive (resp. negative) half-direction. Let R_{ab+} (resp. R_{ab-}) be the vector space on the set C_{ab+} (resp. on C_{ab-}). The decomposition $C_{ab} = C_{ab+} \cup C_{ab-}$ gives $R_{ab+} \oplus R_{ab-} = R_{ab}$. Considering the isomorphism $T_{Z(E)}^{ab} = R_{ab} = R_{ab+} \oplus R_{ab-}$, we will say that R_{ab+} (resp. R_{ab-}) is the positive (resp. negative) tangent space to $\mathbb{H}_{ab}$ at $Z(E)$ and we will write it down $T_{Z(E)}^{ab+}$ (resp. $T_{Z(E)}^{ab-}$). We define $T_{Z(E)}^+ := \oplus T_{Z(E)}^{ab+}$ where the sum runs over all the directions (a, b) .

3 The exponential map

3.1 The invariant case

By the previous section, there is an isomorphism $T_{Z(E)}^{ab+} \simeq \text{Spec } k[X_{c,m}]$ where the $X_{c,m}$ are variables in bijection with the significant cleft couples (c, m) with direction (a, b) and positive half-direction. The goal of this section is to produce a sort of “exponential” map $e : T_{Z(E)}^{ab+} \rightarrow \mathbb{H}_{ab}$ which induces an isomorphism between $T_{Z(E)}^{ab+}$ and the image $C(E)$. In other words, this section provides an explicit chart for the cell $C(E)$. In the remainder of this section, we will make the assumption that the half-direction (a, b) verifies $a > 0$ and $b < 0$.

The morphism $T_{Z(E)}^{ab+} \rightarrow \mathbb{H}$ corresponds to a universal ideal over $\text{Spec } k[X_{c,m}]$ that we describe now. Let $c_1 <_+ c_2 <_+ \dots <_+ c_n$ be the clefts of E . We define a set of monomials Δ_i for $1 \leq i \leq n$ by decreasing induction: Δ_n is the set of monomials divisible by c_n and, for $i \neq n$, Δ_i is the set of monomials divisible by c_i but not divisible by c_{i+1} . We denote by E_i the set of significant cleft couples (c, m) verifying $c = c_i$. We define a polynomial $P(c_i) \in k[X_{c,m}][x, y]$ for each cleft c_i and a polynomial $Q(c_i, m) \in k[X_{c,m}][x, y]$ for each $(c_i, m) \in E_i$ by decreasing induction on i . When $i = n$, we put

$$P(c_n) = c_n.$$

The set E_n being empty, there is no polynomial $Q(c_n, m)$ to define. For a general i and a significant cleft couple (c_i, m) , g the s.c.m. of c_i and c_{i+1} , let c_{i+k} ($k > 0$) be the cleft such that $m.(g/c_i)$ is in Δ_i . We put

$$Q(c_i, m) = P(c_{i+k}). \frac{m}{c_{i+k}}$$

and

$$P(c_i) = P(c_{i+1}). \frac{c_i}{c_{i+1}} + \sum_{m \text{ s.t. } (c_i, m) \text{ is significant}} X_{c_i, m} Q(c_i, m).$$

Proposition 10. a) The ideal I_+ generated by the $P(c_i)$ defines a flat family Z_+ of constant length over $\text{Spec } k[X_{c,m}]$.
b) The fiber of Z_+ over the origin is $Z(E)$.

Proof: when $X_{c_i, m} = 0$, for all (c_i, m) , then $P(c_i) = c_i$ and $I_+ = I^E$ This shows point b).

To verify a) we use the theory of Gröbner bases. It suffices to find a monomial order on $\text{Spec } k[x, y]$ such that, with respect to that monomial order, the $P(c_i)$ are a Gröbner basis of I_+ over each point and verify $\text{in}(P(c_i)) = c_i$. Indeed, if it is the case, the monomials which are in E form a basis of $k[x, y]/I_+$ over each point, so the length of the fibers of $k[x, y]/I_+$ equals the cardinal of E . Each fiber being supported by the origin of $\mathbb{A}^2$, the assertion on the length of the fibers implies the flatness. We choose the monomial order $>_-$. According to the Buchberger algorithm, to verify that the P_i form a Gröbner

basis, we must verify that if c_i and c_{i+1} are two consecutive cleft couples, and if g is the s.c.m. of c_i and c_{i+1} , then the remainder of a division of $h = \frac{g}{c_i}P(c_i) - \frac{g}{c_{i+1}}P(c_{i+1})$ by the $P(c_i)$ is zero. But $h = \frac{g}{c_i} \sum_m$ s.t. (c_i, m) is significant $X_{c_i, m}Q(c_i, m)$. By construction, $\frac{g}{c_i}Q(c_i, m)$ is a product of a polynomial $P(c_{i+k})$ by a monomial. So the remainder of a division is zero. ■

Theorem 11. *The morphism $e : T_{Z(E)}^{ab+} \rightarrow \mathbb{H}$ defined by I_+ induces an isomorphism $T_{Z(E)}^{ab+} \simeq C(E)$.*

Proof: by construction, the image of e verifies $Im(e) \subset C(E)$. If e is an embedding, then $Im(e) = C(E)$ and the theorem is proved. Indeed, suppose that e is an embedding and that there exists $p \in C(E) - Im(e)$. Then the orbit $T_{ab}.p$ is such that $T_{ab}.p \cup Im(e)$ is singular at $Z(E)$. Counting the dimensions of the tangent spaces gives the inequality

$$\dim T_{C(E), Z(E)} \geq \dim T_{T_{ab}.p \cup Im(e), Z(E)} > \dim T_{Z(E)}^{ab+}.$$

This contradicts the equality of the first and last tangent spaces asserted by Bialynicki-Birula.

We will conclude by showing using Plücker coordinates that $i \circ e : T_{Z(E)}^{ab+} \rightarrow \mathbb{P}^N$ is an embedding, where $i : \mathbb{H}^{card(E)} \rightarrow \mathbb{P}^N$ is an embedding of the Hilbert scheme in a projective space.

Fix $n > 0$ and define the set of monomials C_n by

$$C_n := \{m = x^\alpha y^\beta \notin E \text{ such that } d(m) \leq n\}$$

Each m of C_n is in one sector Δ_i . Put $f_m = P(c_i) \cdot \frac{m}{c_i}$. A division of f_m by the elements $f_{m'}$ smaller than f_m with respect to the order $>_-$ is:

$$f_m = \sum_{d(m')=d(m), m' >_+ m} q_{m'} f_{m'} + g_m, \text{ where } q_{m'} \in k.$$

The remainder g_m writes down

$$g_m = m + \sum_{m_i \in E, d(m_i)=d(m)} \mu_{m, m_i} m_i.$$

By construction of $Hilb(\mathbb{A}^2)$, if we have chosen n big enough, $i \circ e$ is given in Plücker coordinates by $\bigwedge_{m \in C_n} g_m$. The coordinate corresponding to the term $\bigwedge_{m \in C_n} m$ equals 1 and $i \circ e$ is then a morphism from $Spec k[X_{c, m}]$ to an affine space $Spec k[Y_1, \dots, Y_p]$. If (c, m) is a significant cleft couple, the coordinate corresponding to the factor $m \bigwedge (\bigwedge_{m' \in C_n, m' \neq c} m')$ is $\mu_{c, m} = X_{c, m} + R_{c, m}$, where $R_{c, m}$ is a polynomial in the variables $X_{c', m'}$, with $c' <_- c$ or $(c' = c \text{ and } m' <_+ m)$. This means that we can order the variables $X_{c, m}$ and the variables $Y_1, \dots, Y_p$ such that the morphism

$$i \circ e : Spec k[X_{c, m}] =: Spec k[X_1, \dots, X_q] \rightarrow Spec k[Y_1, \dots, Y_p]$$

is given by $Y_1 = X_1$ and for $1 < i \leq q$, $Y_i = X_i + R_i$ where R_i is a polynomial in the variables X_j , $j < i$. This shows that $i \circ e$ is an embedding. ■

3.2 The general case

In the previous section, we have defined a map $e : T_{Z(E)}^{ab+} \rightarrow \mathbb{H}_{ab}$ which induces an isomorphism $e : T_{Z(E)}^{ab+} \rightarrow C(E)$. In this section, we extend the map e to a map $T_{Z(E)}^+ \rightarrow \mathbb{H}$, which induces an isomorphism between $T_{Z(E)}^+$ and a Bialynicki-Birula cell $C(E)$ of $\mathbb{H}$ and we explain the link between this description of $C(E)$ and the one of [5]. Note that this section is not useful for the proof of theorem 1.

Choose $a < 0$ and $b < 0$ relatively prime. The one-dimensional torus $k^* = T_{ab}$ acts on $\mathbb{H}$ with the monomial subschemes as the only invariants subschemes by an argument similar to 7, so we can apply Bialynicki-Birula to define cells from this action. To be concrete, the cell $C(E)$ parametrizes the subschemes X such that $\lim_{t \rightarrow 0} (t^{-b}, t^a).X = Z(E)$.

Choose a variable $X_{c,m}$ for each cleft couple with positive half-direction. Define as in the previous section polynomials $P(c_i) \in k[X_{c,m}][x, y]$ for each cleft c_i , $Q(c_i, m) \in k[X_{c,m}][x, y]$ for each significant (c_i, m) . Explicitly, with the notations of the previous section

$$P(c_n) = c_n$$

$$Q(c_i, m) = P(c_{i+k}) \cdot \frac{m}{c_{i+k}}$$

and for $i < n$,

$$P(c_i) = P(c_{i+1}) \cdot \frac{c_i}{c_{i+1}} + \sum_{m \text{ s.t. } (c_i, m) \text{ is significant}} X_{c_i, m} Q(c_i, m).$$

Proposition 12. *The ideal I_+ generated by the $P(c_i)$ defines a flat family Z_+ of constant length over $\text{Spec } k[X_{c,m}]$.*

Proof: this is exactly the same proof as in 10 except for the claim of flatness which is not valid since the family is not any longer supported by the origin. To conclude that Z_+ is flat over $\text{Spec } k[X_{c,m}]$ knowing that the fibers are of constant length, we need to check that if p is the generic point of $\text{Spec } k[X_{c,m}]$, if G_p is the support of the fiber $(Z_+)_p$, then the closure of G_p in $\text{Spec } k[X_{c,m}] \times \mathbb{P}^2$ is a subscheme of $\text{Spec } k[X_{c,m}] \times \mathbb{A}^2$. The ideal of G_p contains an ideal $(x, y^l + a_{l-1}y^{l-1} + \dots a_0y^0)$ where a_i is a polynomial in the indeterminates $X_{c,m}$. The closure $\overline{G_p}$ of G_p in $\text{Spec } k[X_{c,m}] \times \mathbb{P}^2$ is included in the subscheme defined by the homogeneous ideal $(x, y^l + a_{l-1}y^{l-1}h + \dots a_0y^0h^l)$. Thus $\overline{G_p}$ doesn't meet the relative line $h = 0$. ■

Theorem 13. *The morphism $e : T_{Z(E)}^{ab+} \rightarrow \mathbb{H}$ defined by I_+ induces an isomorphism $T_{Z(E)}^{ab+} \simeq C(E)$.*

Proof: the proof is essentially identical to the proof of 11, with few changes left to the reader. ■

Remark 14. *We can clearly by the same method parametrize the cells of the action of a torus T_{ab} , $a > 0, b > 0$ using the negative tangent space.*

We now describe the link between this last theorem and theorem 2 of [5]. In the remainder of this section, we assume familiarity with the notations and the results of [5].

In theorem 2 of [5], Ellingsrud and Strømme consider the Bialynicki-Birula stratification corresponding to an action of a one parameter subgroup $\psi : \mathbb{G} \rightarrow T$. They construct a set of variables s_{ij} . The one dimensional space generated by s_{ij} is provided with an action of the torus $\psi(\mathbb{G})$ of weight n_{ij} . Denoting by $S(\psi)$ the polynomial ring whose indeterminates are the variables s_{ij} of negative weight, they construct a family $Z \subset \text{Spec } S(\psi) \times \mathbb{A}^2$, flat on $\text{Spec } S(\psi)$.

In case $\psi(\mathbb{G})$ coincides with our torus T_{ab} , we can make the link between the two theorems.

In this particular case where $\psi(\mathbb{G}) = T_{ab}$ with $a < 0$ and $b < 0$, the variables s_{ij} with negative weight are the variables s_{ij} with $(i, j) \in \Delta_1$ with their notations. We make a bijection φ between the set R_+ of significant positive cleft couples and Δ_1 . If $(c, m) \in R_+$, let $l > 0$ be the smallest integer such that $x^l \cdot m \notin E$. Using the numeration of the monomials of [5], we have $cx^l = d_j$ and $mx^l = d_i$. We let $\varphi(c, m) = (i, j)$. The following proposition whose proof is omitted says that, up to sign, φ identifies the family Z_+ giving theorem 13 and the family Z giving theorem 2 of Ellingsrud and Strømme.

Proposition 15. *There exists an application $\epsilon : R_+ \rightarrow \{1, -1\}$ such that, if $\eta : \text{Spec } S(\psi) \rightarrow \text{Spec } k[X_{c,m}]$ is the morphism defined by $X_{c,m} \mapsto \epsilon(c, m)s_{\varphi(c,m)}$, then $Z = Z_+ \times_{\text{Spec } k[X_{c,m}]} \text{Spec } S(\psi)$.*

Remark 16. *It seems that the theorem of Ellingsrud and Strømme is more general than ours, since they can take any one parameter subgroup of $k^* \times k^*$ whereas we only consider the sub-tori T_{ab} with $ab > 0$. However, there is a gap in theorem 2 in the extra cases since the family they produce is not always of constant length. For instance, take the one parameter subgroup $t \mapsto (t, t)$, and, following the procedure of [5], construct a family whose fiber over the origin is the subscheme $Z(E)$ of length 6, where $I^E = (y^3, xy, x^4)$. The fiber of Z over the point with all coordinates $s_{ij} = 0$ but $s_{03} = 1, s_{74} = 1$ is of length 7 since a Gröbner basis with respect to the homogeneous order with $x > y$ is $xy^2 + y^3, x^2y + xy^2, x^3 + x^2y - xy - y^2, y^4 - y^3$. In particular, one may see theorem 13 and remark 14 as sufficient conditions under which the theorem of Ellingsrud and Strømme applies.*

Let us explain when these conditions apply. There are 3 types of cells for $\text{Hilb}^l(\mathbb{P}^2)$ in [5], described in theorems 3, 4, 5. We note that the sufficient conditions given in the present paper are exactly the conditions needed in theorem 5. One can show that theorem 3 is correct too (this concerns the cells of $\mathbb{H}$ which are the most important cells). The above counterexample concerns theorem 4.

4 Connecting the strata.

Let $a > 0$ and $b < 0$. The monomials of $k[x, y]$ can be ordered in an infinite sequence $m_0 < m_1 < \dots$ with respect to the order $>$ defined by $m < m'$ if $(d(m), d_y(m)) <$

$(d(m'), d_y(m'))$ for the lexicographic order on $\mathbb{N}^2$. For a staircase E , let S_E be the function from $\mathbb{N}$ to $\mathbb{N}$ defined by $S_E(k)$ = number of monomials in E smaller or equal to m_k .

Definition 17. We define a partial relation $>$ on staircases by: $E > F$ if $\forall k, S_E(k) > S_F(k)$.

The goal of this section is to show the following proposition:

Proposition 18. Let E be a staircase such that $T_{Z(E)}^{ab+} \neq 0$. Then there exists a staircase F verifying $F < E$ and $\overline{C(E)} \cap C(F) \neq \emptyset$.

Proof: choose a point $p \in \text{Spec } k[X_{c,m}]$ different from the origin O . There exists a monomial m which is in $I^E(p)$ but not in $I_+(p)$ (otherwise $I^E(p) \subset I_+(p)$, and even $I^E(p) = I_+(p)$ since the families defined by $I^E(p)$ and $I_+(p)$ have the same relative degree, which is impossible since $e(O) \neq e(p)$ by 11). Take such an m minimal. The element $g_m(p) \in I_+(p)$ defined in theorem 11 has its terms in E except the initial term m . There is at least one term different from m since $m \notin I_+(p)$ by definition. Consider the subscheme $X(p)$ defined by the ideal $I_+(p)$. If the torus k^* acts on $\text{Spec } k[x, y]$ by $t.x = tx, t.y = y$, then the scheme $X(\infty) = \lim_{t \rightarrow \infty} t.X(p)$ is not in $C(E)$ since its ideal contains the monomial $\lim_{t \rightarrow \infty} t.g_m(p) \in E$. If F is the staircase of $X(\infty)$, then $\overline{C(E)} \cap C(F) \neq \emptyset$. This non vacuity implies $E > F$ ([6] or [14]). ■

5 Uniqueness of the final stratum and conclusion of the proof

The connectedness of $\mathbb{H}_{ab}(H)$ has been proved in the first section under the condition $ab \leq 0$. In the remaining cases, one can suppose $a > 0$ and $b < 0$ and we have stratified $\mathbb{H}_{ab}(H)$ by affine spaces $C(E)$, ordered the staircases parametrizing the strata and proven in the previous section that $\overline{C(E)} \cap C(F) \neq \emptyset$ for some staircase $F < E$ provided that $T_{Z(E)}^{ab+} \neq 0$. We can go down from stratum to stratum while the positive tangent space is non trivial. The process stops after a finite number of steps since the number of possible staircases is finite. In fact, the stratum $C(E_m)$ in which the process stops doesn't depend on the intermediate degenerations. The staircase E_m parametrizing this final stratum is characterized by the following theorem.

Recall that a staircase E is compatible with $H = (\dots, h_{-1}, h_0, h_1, \dots)$ if $C(E) \cap H_{ab}(H) \neq \emptyset$ or equivalently if E has h_i elements in degree i .

Theorem 19. Let S be the set of staircases compatible with H . If $S \neq \emptyset$, then there exists in S a staircase E_m such that: $\forall E \in S, E_m \leq E$.

To conclude the proof of the connectedness, it clearly suffices to prove this last theorem and to show that the path through the strata always stops in that special stratum $C(E_m)$.

Since a minimal staircase E verifies $T_{Z(E)}^{ab+} = 0$ by the previous section, and since we

stop when $T_{Z(E)}^{ab+} = 0$, we just have to show that there is a unique staircase $E \in S$ such that $T_{Z(E)}^{ab+} = 0$, which we prove by induction on the cardinal of E .

The case $\text{card}(E) = 1$ is trivial. A general E is the “vertical collision” of the bottom row L and of a residual staircase U determined by its monomials: $E_R := \{m \text{ such that } ym \in E\}$. It is in fact sufficient to show that the bottom row L of E is completely determined. Indeed, suppose that L is known. The positive tangent space at $Z(U)$ injects in the positive tangent space at $Z(E)$ by sending $\sum \lambda_{c,m}(c, m)$ to $\sum \lambda_{c,m}(yc, ym)$, so $T_{Z(U)}^{ab+} = T_{Z(E)}^{ab+} = 0$. It follows that U is completely determined by induction (since its Hilbert function is determined by that of E and L).

The next two lemmas complete the proof. They explain that the bottom row L of a staircase E with trivial positive tangent space and compatible with H is determined: the maximal integer k such that $x^k \in E$ is the maximum integer such that $H(kb - a) < H(kb)$. More precisely, lemma 21 shows that the bottom row contains all the monomials x^k such that $H(kb - a) < H(kb)$ and lemma 20 shows that it cannot contain a bigger monomial.

Lemma 20. *Let E be a staircase compatible with H . Let x^k be the maximal power of x such that $x^k \in E$ and let $\delta = kb$ be its degree. If $H(\delta) \leq H(\delta - a)$, then $T_{Z(E)}^{ab+} \neq \{0\}$.*

Proof: E contains a monomial m of degree δ such that $ym \notin E$. Otherwise, $\varphi : m \mapsto ym$ would be an injective application from the set $E(\delta - a) := \{m \in E, d(m) = b\}$ to $E(\delta)$. The map φ is not surjective since $x^k \notin \text{Im}(\varphi)$. Thus, $\text{card}(E(\delta)) = H(\delta) > H(\delta - a) = \text{card}(E(\delta - a))$, contradicting the hypothesis.

Now choose $l \in \mathbb{N}$ maximal such that $c = \frac{ym}{x^l} \in k[x, y] - E$. By construction, (c, x^{k-l}) is a cleft couple, which shows $T_{Z(E)}^{ab+} \neq 0$. ■

Lemma 21. *Let $\delta = kb$ be an integer. If $H(\delta - a) < H(\delta)$, then for any staircase E compatible with H , the element x^k of degree δ is in the bottom row of E .*

Proof: if x^k were not in E , then to each element $x^\alpha y^\beta \in E$ of E of degree δ would be associated the element $x^{\alpha-1} y^\beta \in E$ of degree $\delta - a$. A count of these elements would then show $H(\delta - a) \geq H(\delta)$. ■

Remark 22. *The existence of a minimal staircase with respect to the partial order on staircases in theorem 19 is a purely combinatorial statement which has required an algebro-geometric proof. I have not found a combinatorial proof simpler than the given argument.*

References

- [1] A Bialynicki-Birula. Some theorems on actions of algebraic groups. *Ann. of Math.*, (98):480–497, 1973.

- [2] J Briançon. Description de $\text{Hilb}^n \mathbb{C}\{X, y\}$. *Invent .Math.*, (41), 1977.
- [3] M Brion. Equivariant chow groups for torus actions. *Transformations Groups*, (2):225–267, 1997.
- [4] Geir Ellingsrud and Stein Arild Strømme. On the homology of the Hilbert scheme of points in the plane. *Invent. math.*, (87):343–352, 1987.
- [5] Geir Ellingsrud and Stein Arild Strømme. On a cell decomposition of the Hilbert scheme of points in the plane. *Invent. Math.*, (91):365–370, 1988.
- [6] L Evain. Incidence relations among the Schubert cells of equivariant Hilbert schemes. *Math. Z.*, to appear.
- [7] J Fogarty. Algebraic families on an algebraic surface. *Am. J. Math.*, (10):511–521, 1968.
- [8] W Fulton. *Introduction to toric varieties*, volume 131. Princeton University Press, 1993.
- [9] R Hartshorne. Connectedness of the Hilbert scheme. *Publ. Math. I.H.E.S.*, (29):5–48, 1966.
- [10] A Iarrobino. Punctual Hilbert schemes. *Mem. Am. Math. Soc.*, (188), 1977.
- [11] B. Iversen. A fixed point formula for action of tori on algebraic varieties. *Inventiones math.*, (16):229–236, 1972.
- [12] M Lehn. Chern classes of tautological sheaves on hilbert schemes of points on surfaces. *Invent. math.*, (136):157–207, 1999.
- [13] H Nakajima. Heisenberg algebra and hilbert schemes of points on projective surfaces. *Ann. of Math.*, 2(145):379–388, 1997.
- [14] J Yameogo. Décomposition cellulaire de variétés paramétrant des idéaux homogènes de $\mathbb{C}[[x, y]]$. *Compositio Math.*, 1(90):81–98, 1994.